
REVIEW

Apoptosis: an introduction for the endodontist

P. G. Satchell¹, J. L. Gutmann² & D. E. Witherspoon³

¹Private Practice Limited to Endodontics, Houston, ²Private Practice Limited to Endodontics, Dallas, and ³Private Practice Limited to Endodontics, Plano, Texas, USA

Abstract

Satchell PG, Gutmann JL, Witherspoon DE. Apoptosis: an introduction for the endodontist. *International Endodontic Journal*, **36**, 237–245, 2003.

Apoptosis plays an important role in many aspects of endodontics, yet there is a paucity of information in this regard in the endodontic literature. Apoptosis is a single deletion of scattered cells by fragmentation into membrane-bound particles that are phagocytosed by other cells. It is a key process in the embryological development of the tooth, periodontal ligament and supporting oral tissue in the progression of oral disease, bone resorption,

immunological response and inflammation, and in wound healing and certain pharmacological effects. The understanding of the ability of clinical materials to induce or inhibit apoptosis and the investigation of apoptosis as it relates to the pathogenesis of pulpal and periradicular pathology may eventually lead to new treatment approaches for the endodontist. The purpose of this review is to familiarize the clinical endodontist with current knowledge on apoptosis as it relates to the pulp and periradicular tissues.

Keywords: apoptosis.

Received 21 May 2001; accepted 2 January 2003

Introduction

Apoptosis is essentially cell suicide, and was first described in the *British Journal of Cancer* in 1972 (Kerr *et al.* 1972). According to these authors, at the appropriate time and under certain conditions, cells self-destruct without damaging adjacent cells. This process is occurring constantly and can happen to seemingly healthy cells. Whilst it is unique to animals and is studied classically in the nematode worm *Caenorhabditis elegans*, it has been increasingly studied *in vitro* and in higher organisms.

There has been an explosion in the number of articles on the subject of apoptosis. A conservative estimate places the number at nearly 45 000. Whilst apoptosis has been a topic of extensive research in the scientific community for some time, it has received little attention in the endodontic literature. The purpose of this article is to familiarise the endodontist with apoptosis and to discuss some of the potential implications of apoptosis to endodontics.

Why and how is apoptosis important to endodontics?

Apoptosis plays a ubiquitous role in the body. Amongst other things, apoptosis is a key process in oral development (Vaahtokari *et al.* 1996, Harada *et al.* 1998, Cerri *et al.* 2000), the progression of oral disease (Polverini & Nor 1999), periradicular lesion development (Onishi *et al.* 1997), resorption (Hughes & Boyce 1997), immunological response and inflammation (Onishi *et al.* 1997), wound healing (Fanning *et al.* 1999) and in certain pharmacological effects (Hughes *et al.* 1995). The understanding of the ability of clinical materials to selectively induce or inhibit apoptosis is leading to new treatment modalities (Bamford *et al.* 2000). An understanding of the mechanisms of apoptosis may lead to adjunctive treatments for pulpal and periradicular diseases through yet unexplored pathways.

What is the apoptotic process?

The entire process of apoptosis takes about 1 h from initiation (Figs 1 and 2). The initiating triggers are many and varied, and are grouped broadly as physiological or nonphysiological. These include, but are not limited to,

Correspondence: Dr James L. Gutmann, 1416 Spenwick Terrace, Dallas, Texas, TX 75204-5529, USA (Tel.: +1 214 827 5378; fax: +1 214 827 4848; e-mail: jlgutmann@earthlink.net).




Figure 1 The major stages of apoptosis include the initiating trigger (1) that leads to the activation of the intracellular mechanism for the apoptotic process (2). Morphological changes of the cell (3) include cell membrane changes that signal phagocytic cell recognition and elimination (4) without promoting inflammation.

the following: Fas ligands (Fas), tumour necrosis factor (TNF), nerve growth factor (NGF), nitric oxide (NO), lipopolysaccharide (LPS), host immune reactions, kinins and glucocorticoids (McKenna *et al.* 1998). The best characterised apoptotic trigger is the Fas ligand, a member of the TNF super-family. The Fas receptor is a cell surface glycoprotein that mediates apoptotic signals from the cell surface into the cytoplasm (Yoshioka *et al.* 1996). When the Fas ligand binds to the Fas receptor on the cell membrane, the newly formed Fas complex is allowed to associate with intracellular proteins. The morphological changes of specific intracellular proteins induced by this complex result in the activation of other substances such as IL-1 β converting enzyme (ICE).

In this particular mechanism, and there are many, Procaspase (inactive form) is activated to the protease Caspase. An amplification cascade then ensues with Caspases activating other Caspases, eventually cleaving the host cell by acting on a variety of cell structures such

as the nuclear membrane (Nicholson & Thornberry 1997). The cell shrinks in the process and there is a loss of cell-cell junctions resulting in detachment from adjacent cells. The chromatin condenses, the cytoplasm 'blebs' (forms so-called 'pseudopods') and the cell breaks up into fragments known as 'apoptotic bodies'. Indirectly activated endonucleases lead to breakdown of the DNA (Kerr *et al.* 1972) into multiples of 180–200 base pair fragments (Cohen & Duke 1984). Finally, either macrophages or adjacent cells phagocytose the apoptotic bodies (Kerr *et al.* 1972) (Figs 1 and 2).

How does necrosis compare to apoptosis?

Cells usually die either by necrosis or apoptosis. The characteristics of apoptotic death are more clearly understood when compared to the characteristics of necrotic death (Table 1). Necrosis is a pathological death of cells resulting from irreversible damage that occurs




Figure 2 Various initiating triggers effect the activation of a central apoptotic signal. This central signal that can be blocked by bcl-2 leads to the activation of proteolytic enzymes via the caspase cascade. Those enzymes lead to morphological changes in the cell structures including the nucleus, cytoskeleton and membrane. Cell membrane changes include expression of proteins for recognition by phagocytic cells.

in the dental pulp and is a term commonly used in pulpal diagnosis. The earliest irreversible changes are mitochondrial, consisting of swelling and granular calcium

Table 1 Comparison of classic features of apoptosis (programmed cell death) and necrosis (pathologic cell death) (Polverini & Nor 1999)

Apoptosis	Necrosis
Controlled process	Uncontrolled process
Energy dependent (ATP required)	No energy required (passive)
Cells shrink (apoptotic bodies)	Cells swell (lysis)
No loss of membrane integrity	Membrane integrity lost
Noninflammatory	Inflammatory
No scarring occurs	Scarring
Individual or small cell groups	Large cell groups (organ segments)
Nuclear fragmentation	Nuclear dissolution
Physiological stimuli	Pathologic stimuli

deposits. After such changes, the outlines of individual cells are indistinct and affected cells may become merged, sometimes forming a focus of coarsely granular, amorphous or hyaline material (Stedman 1995). These features include cell swelling, membrane lysis and an inflammatory response (Wyllie *et al.* 1980), and are distinctly different from the features of apoptosis described above.

What role does apoptosis play in development?

Apoptosis plays an important role in all stages of life. Developing human branchial arches, embryonic tails and finger webbing cannot resorb before birth without organised programmed cell death. In mature organisms, homeostasis is maintained by balancing the continuous

mitosis and differentiation of cells with the apoptotic process (Kerr *et al.* 1972).

Apoptosis has multiple roles in tooth development from the beginning of tooth formation to the completion of root development. There is evidence of apoptosis in the reduction of cells of the stellate reticulum, at the initiation of enamel formation (Vaahtokari *et al.* 1996, Baratella *et al.* 1999) and in the stratum intermedium (Bronckers *et al.* 1996, Vaahtokari *et al.* 1996). This process also occurs during the transition stage between secretion and maturation of ameloblasts during enamel formation (Nishikawa & Sasaki 1995). After enamel formation, approximately 25% of the ameloblasts die and following enamel matrix maturation, another 25% undergo apoptosis (Joseph *et al.* 1994). Apoptosis has been shown to occur around the crowns of teeth during tooth eruption (Bronckers *et al.* 1996, Kaneko *et al.* 1997), playing a major role in the elimination of reduced ameloblasts located at cusps (Shibata *et al.* 1995). It is thought that the inhibition of apoptosis in reduced ameloblasts may occur after the elimination of the tooth organ, allowing for the establishment of the junctional epithelium (Shibata *et al.* 1995). Whilst the precise mechanisms for these events remain largely undiscovered, some work has shown that bcl-2, an apoptotic inhibitor, is involved in maintaining the viability of the enamel organ during tooth development (Slootweg & de Weger 1994).

Fibroblast-like cells of the periodontal ligament (PDL) exhibit apoptosis during tooth development (Cerri *et al.* 2000), and apoptotic cells, probably osteoclasts, are also found on the surfaces of developing alveolar bone (Vaahtokari *et al.* 1996). Apoptosis is also responsible for at least partial elimination of the cells of Hertwig's epithelial root sheath (HERS) after root formation is complete (Kaneko *et al.* 1999, Cerri *et al.* 2000). Furthermore, cell rests of Malassez, which can proliferate to produce periradicular cysts, may be partially as a result of incomplete apoptosis of HERS.

What role does apoptosis play in the dental pulp?

Apoptosis is a part of normal pulp homeostasis (Nishikawa & Sasaki 1999), occurring more in the occlusal (incisal) than in the apical portions of the pulp (Vermelin *et al.* 1996, Nishikawa & Sasaki 1999). Most apoptotic cells in normal pulp can be found at the periphery and are usually associated with the subodontoblastic region rather than with the odontoblastic layer (Vermelin *et al.* 1996, Piattelli *et al.* 2001). Odontoblasts seem to compensate for the reduction in pulp chamber volume as a result of physiological (secondary) dentine

formation by odontoblastic layering rather than by cell death. Apoptosis is more evident in odontoblasts after injury to odontoblastic processes as seen with cavity preparation (Goldberg *et al.* 1994, Bronckers *et al.* 1996, Kitamura *et al.* 2001). There is evidence that reparative (tertiary) dentine formation, in response to injury, is associated with a large decrease in the number of odontoblasts. Up to half of the pulp odontoblasts can be eliminated in only 4 years by this process (Franquin *et al.* 1998). Fibroblasts and vascular endothelial cells of the pulp proper also show evidence of apoptosis (Franquin *et al.* 1998). Although certain characteristics of apoptotic death are favourable to necrosis (i.e. an inflammatory response is not triggered), apoptosis does not lead to pulp recovery. This may be due in part to the elimination of regenerative cells by apoptosis (Goldberg *et al.* 1994). Ultimately, when apoptosis occurs, apoptotic bodies and debris are phagocytosed by major histocompatibility Class II (MHC II) positive and negative macrophages (Nishikawa & Sasaki 1999).

What role does apoptosis play in other oral tissues?

Apoptosis seems to be necessary for maintaining homeostasis within continually renewing tissues such as the oral mucosa and skin (Funato *et al.* 1999). Gingival tissue has a high cell turnover, and apoptosis has been demonstrated to occur in this tissue in 90% of the individuals tested (Yoshioka *et al.* 1996). Here, as in other parts of the body, apoptosis is essentially a counterbalance to mitosis. Unfortunately, the role of apoptosis in the differentiation of oral epithelial cells is not clear (Harada *et al.* 1998), and more work is required to fully describe the events associated with this process in normal oral tissues as a baseline to further study.

What role does apoptosis play in bone?

In the process of bone remodelling, some osteoblasts die via apoptotic mechanisms, whilst those remaining become embedded as osteocytes (Ihbe *et al.* 1998, Jilka *et al.* 1998). Many osteoclasts that lose their attachment to bone die by the process of apoptosis (Hughes & Boyce 1997). In this regard, a third term in addition to necrosis and apoptosis is oncosis, which describes cell death associated with slow ischaemia and cell swelling. This occurs to some osteoblasts during transition to osteocytes (Darzynkiewicz & Traganos 1998).

TNF- α has been shown to enhance osteoblast apoptosis and may contribute to bone loss associated with inflammation (Hill *et al.* 1997, Jilka *et al.* 1998, Tsuboi

et al. 1999). Neutrophils can also induce apoptosis of osteoblasts, demonstrating the relationship between inflammatory cells and bone resorption associated with inflammation (Kawakami *et al.* 1997). On the other hand, growth factors, cytokines and other bone-stimulating hormones can reduce apoptosis of osteoblasts during periods of bone resorption (Hill *et al.* 1997, Jilka *et al.* 1998).

Apoptosis occurs in periradicular tissue during bone remodelling associated with orthodontic tooth movement (Rana *et al.* 2001). Certain mediators may limit the resorptive process and aid in bone formation during remodelling (Hill *et al.* 1997). There is significant evidence that osteoblasts may be involved in the regulation of osteoclast apoptosis (Fuller *et al.* 1993; 1998, Greenfield *et al.* 1999). In fact, most of the mediators that stimulate osteoclast activity seem to act through osteoblasts (Greenfield *et al.* 1999). In general, however, factors that stimulate bone resorption inhibit osteoclast apoptosis and factors that inhibit bone resorption promote osteoclast apoptosis (Hughes & Boyce 1997).

Apoptosis of osteoclast precursors may be one way that the osteoclast cell population is controlled (Hughes & Boyce 1997), effectively reducing bone resorption (van't Hof & Ralston 1997). Furthermore, members of the TNF super-family inhibit osteoclast apoptosis (Fuller *et al.* 1998) and induce precursor maturation to mature functioning osteoclasts (Fuller *et al.* 1998), thereby contributing to bone loss.

In the presence of high extracellular calcium concentrations as a result of ongoing resorptive processes, osteoclast apoptosis is induced (Lorget *et al.* 2000). Similarly, oestrogen, glucocorticoids, bisphosphonates and TGF- β have been shown to stimulate apoptosis of osteoclasts (Greenfield *et al.* 1999).

What role does apoptosis play in immune cells?

Apoptosis is associated with the maintenance of immune cell homeostasis (Usherwood *et al.* 1999). Neutrophil production is balanced by apoptosis and clearance from tissues without inducing an inflammatory response (Onishi *et al.* 1997). Apoptosis also plays a critical role in eliminating harmful or injured cells from tissues. This suggests its participation in inflammatory processes and in the resolution of inflammatory reactions (Onishi *et al.* 1997). In periradicular lesions, apoptosis occurs predominantly in neutrophils (Takahashi *et al.* 1999), a process that may be defective in abscess formation as a result of the acidic environment (Onishi *et al.* 1997). This lack of clearance of apoptotic cells can present a persistent antigen, inducing an autoimmune

response (Onishi *et al.* 1997). Apoptosis of neutrophils is the major means of ending neutrophil-associated inflammation (Marshall & Watson 1997). Induction of this process therefore could be a way of minimizing inflammation that would occur as a result of neutrophil necrosis (Onishi *et al.* 1997), which is mediated and induced by certain environmental signals (Marshall & Watson 1997). There is also evidence that the failure of apoptotic mechanisms in neutrophils can contribute to pathogenesis of disease (Marshall & Watson 1997). Neutrophils in exudate show delayed apoptosis and loss of TNF- α , which is necessary for neutrophil survival in the extravascular fluid (Seely *et al.* 1998). In addition, neutrophils in diabetics do not undergo lipopolysaccharide (LPS)-induced inhibition of apoptosis. This may contribute to the increased susceptibility to infection observed in diabetic patients (Tennenberg *et al.* 1999). Calcium ions have been shown to affect neutrophil apoptosis (Onishi *et al.* 1997). The signalling mechanisms for other immune cells have been studied less extensively than those for neutrophils.

What is the role of apoptosis in oral disease?

Apoptosis is widely involved in disease mechanisms of the oral cavity. Oral diseases in which apoptosis plays a role include lichen planus (Dekker *et al.* 1997, Bloor *et al.* 1999), odontogenic keratocysts (Muraki *et al.* 1997), leucoplakia (Muraki *et al.* 1997), squamous cell carcinoma (Muraki *et al.* 1997, Kaur & Ralhan 2000, Ravindranath *et al.* 2000), oral lymphoma (Regezi *et al.* 1998), aphthous ulceration (Honma *et al.* 1985), erythema multiforme (Chrysomali *et al.* 1997), Sjögren's syndrome (Ishimaru *et al.* 2000) and mucocutaneous candidiasis (Heidenreich *et al.* 1996).

Can microorganisms induce or inhibit apoptosis?

Bacterial proteases have been implicated as inducers of host cell apoptosis (Wang *et al.* 1998). *Fusobacterium nucleatum* induces apoptosis in peripheral blood mononuclear cells and neutrophils (Jewett *et al.* 2000). This is significant because premature immune cell apoptosis may decrease host resistance and promote infection (Geatch *et al.* 1999).

Furthermore, apoptosis of both neutrophils and monocytes decreases in the presence of LPS (Mangan *et al.* 1993, Preshaw *et al.* 1999, Tennenberg *et al.* 1999), resulting in a heightened immune response (Hiroi *et al.* 1998). Another organism, *Prevotella intermedia*, does not induce apoptosis in cultured human osteoblastic

cells (Morimoto *et al.* 1999), whilst in *Streptococcus mutans*, lipoteichoic acid may be a factor in pulpitis as there is evidence that it causes apoptosis in human dental pulp cells of deciduous teeth (Wang *et al.* 2001). Generally, these interactions are complex and unpredictable as microorganisms may increase apoptosis in some host cells and decrease apoptosis in others.

Whilst bacteria and their by-products can induce pulpal inflammation, they do not always lead to pulp necrosis (Bergenholtz 2000). Bacterial proteases have been implicated as inducers of host cell apoptosis (Wang *et al.* 1998). Although certain characteristics of apoptotic death are favourable to necrosis, apoptosis does not lead to pulpal regeneration (Goldberg *et al.* 1994).

How is apoptosis involved in periodontal disease?

There is a considerable amount of research on the role of apoptosis in periodontal disease. Apoptosis occurs in cells of the periodontium as part of normal turnover and remodelling (Koulouri *et al.* 1999), and may be even more prevalent than necrosis in periodontal disease (Sorkin & Niederman 1998). It seems to play a role in age regulation of some immune cells and may be involved in the maintenance of local immune homeostasis in inflamed gingival tissue (Tonetti *et al.* 1998). There is also evidence that increased inflammation is associated with increased epithelial cell apoptosis in the periodontium of patients with periodontal disease (Carro *et al.* 1997).

Various microorganisms associated with periodontal diseases have been shown to generate different short-chain carboxylic acids as metabolic by-products, which can promote inflammation by inhibiting normal apoptosis of certain inflammatory cells (Niederman *et al.* 1997). For example, lactic acid and propionic acid commonly produced by oral microorganisms in periodontal diseases have been shown to inhibit apoptosis of various inflammatory cells (Niederman *et al.* 1997, Yamamoto *et al.* 1997, Sorkin & Niederman 1998). The resultant presence of increased numbers of inflammatory cells results in an elevated inflammatory state.

What is the relationship between gingival trauma, wound healing and apoptosis?

There is a strong evidence that trauma, including surgical trauma, inhibits immune cell apoptosis (Ertel *et al.* 1999, Fanning *et al.* 1999, Ogura *et al.* 1999, Nolan *et al.* 2000). Apoptosis occurs in cells at the advancing epithelial edge in wound healing (Brown *et al.* 1997). There is speculation that the signal for apoptosis and downregulation

of inflammation in a wound may in fact be derived from the epithelium because it appears concurrently with re-epithelialisation of the wound (Brown *et al.* 1997, Leonardi *et al.* 2001).

Granulation tissue fibroblasts (myofibroblasts) play a role in wound contraction. When granulation tissue evolves into a scar, myofibroblasts disappear, probably as a result of apoptosis. Myofibroblasts persist in excessive scarring conditions (Desmouliere 1995), possibly because of an inappropriate inhibition of apoptosis in these cells. Certain mediators may be potential stimulators of apoptosis in myofibroblasts after re-epithelialization in the palatal wound healing process (Funato *et al.* 1999).

What is the role of apoptosis in pharmacotherapeutics?

The principle therapeutic goal of apoptosis research is the understanding of how to induce or inhibit apoptosis in specific cells. This is probably the best approach for altering rates of apoptosis in target cells whilst avoiding systemic toxic effects that may be otherwise associated. Aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), and selective cyclooxygenase 2 (COX2) inhibitors induce apoptosis in certain cells via a caspase activation mechanism. Certain drug combinations may induce apoptosis by a mechanism involving the disruption of the cell cycle (Bamford *et al.* 2000).

The induction of osteoclast apoptosis for the treatment of bone resorption is another therapy that utilizes apoptotic mechanisms (Hughes & Boyce 1997). Antiresorptive drugs such as the bisphosphonate family have been shown to inhibit bone resorption by inducing osteoclast (Hughes *et al.* 1995, Rogers *et al.* 1996, Hiroi-Furuya *et al.* 1999) and macrophage apoptosis (Rogers *et al.* 1996). Bisphosphonates inhibit osteocyte and osteoblast apoptosis contributing to their antiresorptive effects (Plotkin *et al.* 1999). Tetracycline can also inhibit resorption by inducing osteoclast apoptosis (Cillari *et al.* 1998, Vernillo & Rifkin 1998). This is part of the process by which doxycycline downregulates the inflammatory process (Liu *et al.* 1999). There is some thought that tetracycline uses a unique mechanism to induce this selective apoptosis (Bettany & Wolowacz 1998); however, details of this mechanism have not been fully described.

Which dental materials have apoptotic effects?

An understanding of the difference between necrotic and apoptotic forms of cell death is important for understanding the pulp reaction to dental materials (Goldberg

et al. 1994). In cultured cells derived from the human PDL, evidence of apoptosis in response to different dental materials has been described (Adams et al. 1995) and may occur in damaged areas of the pulp because of certain toxic substances in those materials (Goldberg et al. 1994). In some cases, dental materials may lead to uncontrolled development of apoptosis within central pulp cells, but may leave the odontoblasts more or less intact (Goldberg et al. 1994).

More specifically, and related to endodontic treatment, *in vitro* apoptotic changes are documented in human PDL cells exposed to calcium hydroxide (Adams et al. 1995). Calcium hydroxide has antibacterial properties and has shown predictable healing and hard tissue formation when used in cavity preparations (Bergenholtz 2000). Calcium ions may also affect neutrophil apoptosis (Onishi et al. 1997). Unfortunately, neither the apoptotic effects of composite resin on the dental pulp nor the potential apoptotic effects of extruded root canal sealers on the cells that make up the supporting periradicular tissues have been fully documented. Therefore, the long-term effects of these treatment variables may or may not influence the ultimate treatment outcome and should not be overlooked as potential aetiological factors in failure.

References

- Adams AM, Soames JV, Searle RF (1995) Ultrastructural changes in cultured human periodontal ligament cells exposed to dental materials. *Journal of Biomedical Materials Research* **29**, 999–1004.
- Bamford M, Walkinshaw G, Brown R (2000) Therapeutic applications of apoptosis research. *Experimental Cell Research* **256**, 1–11.
- Baratella L, Arana-Chavez VE, Katchburian E (1999) Apoptosis in the early involuting stellate reticulum of rat molar tooth germs. *Anatomy and Embryology* **200**, 49–54.
- Bergenholtz G (2000) Evidence for bacterial causation of adverse pulpal responses in resin-based dental restorations. *Critical Reviews of Oral Biology and Medicine* **11**, 467–80.
- Bettany JT, Wolowacz RG (1998) Tetracycline derivatives induce apoptosis selectively in cultured monocytes and macrophages but not in mesenchymal cells. *Advances in Dental Research* **12**, 136–43.
- Bloor BK, Malik FK, Odell EW, Morgan PR (1999) Quantitative assessment of apoptosis in oral lichen planus. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **88**, 187–95.
- Bronckers AL, Lyaruu DM, Goei Wet al. (1996) Nuclear DNA fragmentation during postnatal tooth development of mouse and hamster and during dentin repair in the rat. *European Journal of Oral Sciences* **104**, 102–11.
- Brown DL, Kao WW, Greenhalgh DG (1997) Apoptosis down-regulates inflammation under the advancing epithelial wound edge: delayed patterns in diabetes and improvement with topical growth factors. *Surgery* **121**, 372–80.
- Carro OM, Evans SA, Leone CW (1997) Effect of inflammation on the proliferation of human gingival epithelial cells *in vitro*. *Journal of Periodontology* **68**, 1070–5.
- Cerri PS, Freymuller E, Katchburian E (2000) Apoptosis in the early developing periodontium of rat molars. *Anatomical Record* **258**, 136–44.
- Chrysomali E, Lozada-Nur F, Dekker NP, Papanicolaou SI, Regezi JA (1997) Apoptosis in oral erythema multiforme. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **83**, 272–80.
- Cillari E, Milano S, D'Agostino P et al. (1998) Modulation of nitric oxide production by tetracyclines and chemically modified tetracyclines. *Advances in Dental Research* **12**, 126–30.
- Cohen JJ, Duke RC (1984) Glucocorticoid activation of a calcium-dependent endonuclease in thymocyte nuclei leads to cell death. *Journal of Immunology* **132**, 38–42.
- Darzynkiewicz Z, Traganos F (1998) Measurement of apoptosis. In: Scheper T, ed. *Advances in Biochemical Engineering/Biotechnology*. Berlin: Springer-Verlag, pp. 33–73.
- Dekker NP, Lozada-Nur F, Lagenaur LA, MacPhail LA, Bloom CY, Regezi JA (1997) Apoptosis-associated markers in oral lichen planus. *Journal of Oral Pathology and Medicine* **26**, 170–5.
- Desmouliere A (1995) Factors influencing myofibroblast differentiation during wound healing and fibrosis. *Cell Biology International* **19**, 471–6.
- Ertel W, Keel M, Buergi U, Hartung T, Imhof HG, Trentz O (1999) Granulocyte colony-stimulating factor inhibits neutrophil apoptosis at the local site after severe head and thoracic injury. *Journal of Trauma-Injury Infection and Critical Care* **46**, 784–93.
- Fanning NE, Porter J, Shorten GD et al. (1999) Inhibition of neutrophil apoptosis after elective surgery. *Surgery* **126**, 527–34.
- Franquin JC, Remusat M, Abou Hashieh I, Dejou J (1998) Immunocytochemical detection of apoptosis in human odontoblasts. *European Journal of Oral Sciences* **106** (Suppl. 1), 384–7.
- Fuller K, Owens JM, Jagger CJ, Wilson A, Moss R, Chambers TJ (1993) Macrophage colony-stimulating factor stimulates survival and chemotactic behavior in isolated osteoclasts. *Journal of Experimental Medicine* **178**, 1733–44.
- Fuller K, Wong B, Fox S, Choi Y, Chambers TJ (1998) TRANCE is necessary and sufficient for osteoblast-mediated activation of bone resorption in osteoclasts. *Journal of Experimental Medicine* **188**, 997–1001.
- Funato N, Moriyama K, Baba Y, Kuroda T (1999) Evidence for apoptosis induction in myofibroblasts during palatal mucoperiosteal repair. *Journal of Dental Research* **78**, 1511–7.
- Geatch DR, Harris JI, Heasman PA, Taylor JJ (1999) *In vitro* studies of lymphocyte apoptosis induced by the periodontal pathogen *Porphyromonas gingivalis*. *Journal of Periodontal Research* **34**, 70–8.

- Goldberg M, Lasfargues JJ, Legrand JM (1994) Clinical testing of dental materials – histological considerations. *Journal of Dentistry* **22** (Suppl. 2), S25–8.
- Greenfield EM, Bi Y, Miyauchi A (1999) Regulation of osteoclast activity. *Life Sciences* **65**, 1087–102.
- Harada H, Mitsuyasu T, Seta Y, Maruoka Y, Toyoshima K, Yasumoto S (1998) Overexpression of bcl-2 protein inhibits terminal differentiation of oral keratinocytes *in vitro*. *Journal of Oral Pathology and Medicine* **27**, 11–7.
- Heidenreich S, Otte B, Lang D, Schmidt M (1996) Infection by *Candida albicans* inhibits apoptosis of human monocytes and monocytic U937 cells. *Journal of Leukocyte Biology* **60**, 737–43.
- Hill PA, Tumber A, Meikle MC (1997) Multiple extracellular signals promote osteoblast survival and apoptosis. *Endocrinology* **138**, 3849–58.
- Hiroi M, Shimojima T, Kashimata M *et al.* (1998) Inhibition by *Porphyromonas gingivalis* LPS of apoptosis induction in human peripheral blood polymorphonuclear leukocytes. *Anticancer Research* **18**, 3475–9.
- Hiroi-Furuya E, Kameda T, Hiura K *et al.* (1999) Etidronate (EHDP) inhibits osteoclastic-bone resorption, promotes apoptosis and disrupts actin rings in isolate-mature osteoclasts. *Calcified Tissue International* **64**, 219–23.
- van't Hof RJ, Ralston SH (1997) Cytokine-induced nitric oxide inhibits bone resorption by inducing apoptosis of osteoclast progenitors and suppressing osteoclast activity. *Journal of Bone and Mineral Research* **12**, 1797–804.
- Honma T, Saito T, Fujioka Y (1985) Possible role of apoptotic cells of the oral epithelium in the pathogenesis of aphthous ulceration. *Oral Surgery, Oral Medicine, Oral Pathology* **59**, 379–87.
- Hughes DE, Boyce BF (1997) Apoptosis in bone physiology and disease. *Molecular Pathology* **50**, 132–7.
- Hughes DE, Wright KR, Uy HL *et al.* (1995) Bisphosphonates promote apoptosis in murine osteoclasts *in vitro* and *in vivo*. *Journal of Bone and Mineral Research* **10**, 1478–87.
- Ihbe A, Baumann G, Heinzmann U, Atkinson MJ (1998) Loss of the differentiated phenotype precedes apoptosis of ROS 17/2.8 osteoblast-like cells. *Calcified Tissue International* **63**, 208–13.
- Ishimaru N, Yoneda T, Saegusa K *et al.* (2000) Severe destructive autoimmune lesions with ageing in murine Sjogren's syndrome through Fas-mediated apoptosis. *American Journal of Pathology* **156**, 1557–64.
- Jewett A, Hume WR, Le H *et al.* (2000) Induction of apoptotic cell death in peripheral blood mononuclear and polymorphonuclear cells by an oral bacterium, *Fusobacterium nucleatum*. *Infection and Immunity* **68**, 1893–8.
- Jilka RL, Weinstein RS, Bellido T, Parfitt AM, Manolagas SC (1998) Osteoblast programmed cell death (apoptosis): modulation by growth factors and cytokines. *Journal of Bone and Mineral Research* **13**, 793–802.
- Joseph BK, Gobe GC, Savage NW, Young WG (1994) Expression and localization of sulphated glycoprotein-2 mRNA in the rat incisor tooth ameloblasts: relationships with apoptosis. *International Journal of Experimental Pathology* **75**, 313–20.
- Kaneko H, Hashimoto S, Enokiya Y, Ogiuchi H, Shimono M (1999) Cell proliferation and death of Hertwig's epithelial root sheath in the rat. *Cell and Tissue Research* **298**, 95–103.
- Kaneko H, Ogiuchi H, Shimono M (1997) Cell death during tooth eruption in the rat: surrounding tissues of the crown. *Anatomy and Embryology* **195**, 427–34.
- Kaur J, Ralhan R (2000) Induction of apoptosis by abrogation of HSP70 expression in human oral cancer cells. *International Journal of Cancer* **85**, 1–5.
- Kawakami A, Eguchi K, Matsuoka N *et al.* (1997) Fas and Fas ligand interaction is necessary for human osteoblast apoptosis. *Journal of Bone and Mineral Research* **12**, 1637–46.
- Kerr JF, Wyllie AH, Currie AR (1972) Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *British Journal of Cancer* **26**, 239–57.
- Kitamura C, Kumra K, Nakayama T, Toyoshima K, Terashita M (2001) Primary and secondary induction of apoptosis in odontoblasts after cavity preparation of rat molars. *Journal of Dental Research* **80**, 1530–4.
- Koulouri O, Lappin DE, Radvar M, Kinane DF (1999) Cell division, synthetic capacity and apoptosis in periodontal lesions analysed by *in situ* hybridisation and immunohistochemistry. *Journal of Clinical Periodontology* **26**, 552–9.
- Leonardi R, Villari L, Caltabiano M, Trivali S (2001) Heat shock protein 27 expression in the epithelium of periapical lesions. *Journal of Endodontics* **27**, 89–92.
- Liu J, Kuszynski CA, Baxter BT (1999) Doxycycline induces Fas/Fas ligand-mediated apoptosis in Jurkat T lymphocytes. *Biochemical and Biophysical Research Communications* **260**, 562–7.
- Lorget F, Kamel S, Mentaverri R *et al.* (2000) High extracellular calcium concentrations directly stimulate osteoclast apoptosis. *Biochemical and Biophysical Research Communications* **268**, 899–903.
- Mangan DE, Mergenhagen SE, Wahl SM (1993) Apoptosis in human monocytes: possible role in chronic inflammatory diseases. *Journal of Periodontology* **64**, 461–6.
- Marshall JC, Watson RW (1997) Programmed cell death (apoptosis) and the resolution of systemic inflammation. *Canadian Journal of Surgery* **40**, 169–74.
- McKenna SL, McGowan AJ, Cotter TG (1998) Molecular mechanisms of programmed cell death. In: Schepers T, ed. *Advances in Biochemical Engineering/Biotechnology*. Berlin Heidelberg: Springer-Verlag, pp. 1–31.
- Morimoto Y, Morimoto H, Murata T, Kobayashi S, Ohba T, Haneji T (1999) Extracts of *Actinobacillus actinomycetemcomitans* induce apoptotic cell death in human osteoblastic MG63 cells. *Journal of Dental Research* **78**, 735–42.
- Muraki Y, Yoshioka C, Fukuda J, Haneji T, Kobayashi N (1997) Immunohistochemical detection of Fas antigen in oral epithelia. *Journal of Oral Pathology and Medicine* **26**, 57–62.
- Nicholson DW, Thornberry NA (1997) Caspases: killer proteases. *Trends in Biochemical Sciences* **22**, 299–306.
- Niederman R, Zhang J, Kashket S (1997) Short-chain carboxylic acid-stimulated, PMN-mediated gingival inflammation. *Critical Reviews in Oral Biology and Medicine* **8**, 269–90.

- Nishikawa S, Sasaki F (1995) DNA localization in nuclear fragments of apoptotic ameloblasts using anti-DNA immunoelectron microscopy: programmed cell death of ameloblasts. *Histochemistry and Cell Biology* **104**, 151–9.
- Nishikawa S, Sasaki F (1999) Apoptosis of dental pulp cells and their elimination by macrophages and MHC class II-expressing dendritic cells. *Journal of Histochemistry and Cytochemistry* **47**, 303–12.
- Nolan B, Collette H, Baker S et al. (2000) Inhibition of neutrophil apoptosis after severe trauma is NF κ B dependent. *Journal of Trauma-Injury Infection and Critical Care* **48**, 599–604 (discussion 604–5).
- Ogura H, Tanaka H, Koh T et al. (1999) Priming, second-hit priming, and apoptosis in leukocytes from trauma patients. *Journal of Trauma-Injury Infection and Critical Care* **46**, 774–81 (discussion 781–3).
- Onishi Y, Tanimoto Y, Kizaki H (1997) Inflammation and apoptosis. *Bulletin of Tokyo Dental College* **38**, 65–76.
- Piattelli A, Rubini C, Fioroni M, Ciavarella L, De Fazio P (2001) Bcl-2, 53, and MIB-1 in human adult dental pulp. *Journal of Endodontics* **26**, 225–7.
- Plotkin LI, Weinstein RS, Parfitt AM, Roberson PK, Manolagas SC, Bellido T (1999) Prevention of osteocyte and osteoblast apoptosis by bisphosphonates and calcitonin. *Journal of Clinical Investigation* **104**, 1363–74.
- Polverini PJ, Nor JE (1999) Apoptosis and predisposition to oral cancer. *Critical Reviews in Oral Biology and Medicine* **10**, 139–52.
- Preshaw PM, Schifferle RE, Walters JD (1999) *Porphyromonas gingivalis* lipopolysaccharide delays human polymorphonuclear leukocyte apoptosis *in vitro*. *Journal of Periodontal Research* **34**, 197–202.
- Rana MW, Pothisiri V, Killiany DM, Xu XM (2001) Detection of apoptosis during orthodontic tooth movement in rats. *American Journal of Dentofacial Orthopedics* **119**, 516–21.
- Ravindranath NM, Nishimoto K, Chu K, Shuler C (2000) Cell-surface expression of complement restriction factors and sialyl Lewis antigens in oral carcinoma: relevance to chemo-immunotherapy. *Anticancer Research* **20**, 21–6.
- Regezi JA, McMillan A, Dekker N et al. (1998) Apoptosis-associated proteins in oral lymphomas from HIV-positive patients. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **86**, 196–202.
- Rogers MJ, Chilton KM, Coxon FP et al. (1996) Bisphosphonates induce apoptosis in mouse macrophage-like cells *in vitro* by a nitric oxide-independent mechanism. *Journal of Bone and Mineral Research* **11**, 1482–91.
- Seely AJ, Swartz DE, Giannias B, Christou NV (1998) Reduction in neutrophil cell surface expression of tumor necrosis factor receptors but not Fas after transmigration: implications for the regulation of neutrophil apoptosis. *Archives of Surgery* **133**, 1305–10.
- Shibata S, Suzuki S, Tengan T, Yamashita Y (1995) A histochemical study of apoptosis in the reduced ameloblasts of erupting mouse molars. *Archives of Oral Biology* **40**, 677–80.
- Slootweg PJ, de Weger RA (1994) Immunohistochemical demonstration of bcl-2 protein in human tooth germs. *Archives of Oral Biology* **39**, 545–50.
- Sorkin BC, Niederman R (1998) Short chain carboxylic acids decrease human gingival keratinocyte proliferation and increase apoptosis and necrosis. *Journal of Clinical Periodontology* **25**, 311–5.
- Stedman TL (1995) *Stedman's Medical Dictionary*, 26th edn. Baltimore, USA: Williams & Wilkins.
- Takahashi K, MacDonald D, Murayama Y, Kinane D (1999) Cell synthesis, proliferation and apoptosis in human dental periapical lesions analyzed by *in situ* hybridization and immunohistochemistry. *Oral Diseases* **5**, 313–20.
- Tennenberg SD, Finkenaue R, Dwivedi A (1999) Absence of lipopolysaccharide-induced inhibition of neutrophil apoptosis in patients with diabetes. *Archives of Surgery* **134**, 1229–33 (discussion 1233–4).
- Tonetti MS, Cortellini D, Lang NP (1998) *In situ* detection of apoptosis at sites of chronic bacterially induced inflammation in human gingiva. *Infection and Immunity* **66**, 5190–5.
- Tsuboi M, Kawakami A, Nakashima T et al. (1999) Tumor necrosis factor- α and interleukin-1 β increase the Fas-mediated apoptosis of human osteoblasts (see comments). *Journal of Laboratory and Clinical Medicine* **134**, 222–31.
- Usherwood EJ, Crowther G, Woodland DL (1999) Apoptotic cells are generated at every division of *in vitro* cultured T-cell lines. *Cellular Immunology* **196**, 131–7.
- Vahtokari A, Aberg T, Thesleff I (1996) Apoptosis in the developing tooth: association with an embryonic signaling center and suppression by EGF and FGF-4. *Development* **122**, 121–9.
- Vermelin L, Lecolle S, Septier D, Lasfargues JJ, Goldberg M (1996) Apoptosis in human and rat dental pulp. *European Journal of Oral Sciences* **104**, 547–53.
- Vernillo AT, Rifkin BR (1998) Effects of tetracyclines on bone metabolism. *Advances in Dental Research* **12**, 56–62.
- Wang PL, Sato K, Oido M et al. (1998) Involvement of CD14 on human gingival fibroblasts in *Porphyromonas gingivalis* lipopolysaccharide-mediated interleukin-6 secretion. *Archives of Oral Biology* **43**, 687–94.
- Wang PL, Shirasu S, Daito M, Ohura K (2001) *Streptococcus mutans* lipoteichoic acid-induced apoptosis in cultured dental pulp cells from human deciduous teeth. *Biochemical and Biophysical Research Communications* **281**, 957–61.
- Wyllie AH, Kerr JF, Currie AR (1980) Cell death: the significance of apoptosis. *International Review of Cytology* **68**, 251–306.
- Yamamoto M, Fujihashi K, Hiroi T, McGhee JR, Van Dyke TE, Kiyono H (1997) Molecular and cellular mechanisms for periodontal diseases: role of Th1 and Th2 type cytokines in induction of mucosal inflammation. *Journal of Periodontal Research* **32**, 115–9.
- Yoshioka C, Muraki Y, Fukuda J, Haneji T, Kobayashi N (1996) Identification of the Fas antigen in human gingiva. *Journal of Dental Research* **75**, 1353–7.